arXiv:0805.1311v1 [math-ph] 9 May 2008

Upper bounds for Total Cross Section in scattering by an obstacle with impedance boundary conditions

J.P.Cruz* Lakshtanov E.L.*†

Keywords: Upper bound, Total Cross Section, Scattering by obstacle, impedance boundary conditions.

Abstract

The scalar scattering of a plane wave by a smooth obstacle with impedance boundary conditions is considered. Upper bounds for the Total Cross Section and for the absorbed power are presented.

---

*Department of Mathematics, Aveiro University, Aveiro 3810, Portugal. This work was supported by *Centre for Research on Optimization and Control* (CEOC) from the "*Fundação para a Ciência e a Tecnologia*" (FCT), cofinanced by the European Community Fund FEDER/POCTI, and by the FCT research project PTDC/MAT/72840/2006.

†e-mail: lakshtanov@rambler.ru

1 Introduction

Consider a bounded body $\Omega \subset \mathbb{R}^3$ with Lipschitz boundary $\partial\Omega$ and $k > 0$. The scattered field is given by the Helmholtz equation and a radiation condition

$$\Delta u(r) + k^2 u(r) = 0, \quad r \in \Omega' = \mathbb{R}^3 \backslash \Omega \tag{1}$$

$$\int_{|r|=R} \left| \frac{\partial u(r)}{\partial |r|} - iku(r) \right|^2 dS = o(1), \quad R \to \infty, \tag{2}$$

with impedance boundary conditions of the form

$$\left(\frac{\partial}{\partial n} + (a + ib) \right) (u(r) + e^{ikz}) = 0, \qquad r = (x, y, z) \in \partial\Omega, \quad a, b \in \mathbb{R}, b > 0 \tag{3}$$

$e^{ikz} = e^{ik(r \cdot \theta_0)}$ is an incident field formed by a plane wave with incident angle $\theta_0 = (0, 0, 1) \in S^2$. In [6], for example, is proved the existence and uniqueness of the solution of (1)-(3). A function $u(r)$ which satisfies the mentioned conditions has asymptotic

$$u(r) = \frac{e^{ik|r|}}{|r|} f_{a,b}(\theta) + o\left(\frac{1}{|r|} \right), \quad r \to \infty, \quad \theta = r/|r| \in S^2, \tag{4}$$

where the function $f_{a,b}(\theta) = f_{a,b}(\theta, k)$ is called *scattering amplitude* and the quantity

$$\sigma_{a,b} = \int_{S^2} |f_{a,b}(\theta)|^2 d\sigma(\theta)$$

is called the Total Cross Section. σ is a square element of the unit sphere.

The power absorbed by the obstacle is given by

$$\mathcal{L} = b\|e^{ikz} + u\|^2_{L_2(\partial\Omega,dS)}$$

It is valid the optical theorem (for example see [2])

$$\sigma_{a,b} = \frac{4\pi}{k} Im[f_{a,b}(\theta_0)] + \frac{1}{k}\mathcal{L}.$$

Theorem 1. *The following inequalities hold for total cross section*

$$\sigma_{a,b} \le \frac{S(k+\sqrt{a^2+b^2})^2(b+\sqrt{a^2+b^2})}{kb^2} \tag{5}$$

and for the power absorbed by the obstacle

$$\mathcal{L} \le \frac{S(k+b+\sqrt{a^2+b^2})^2}{b}. \tag{6}$$

Note that similar inequalities for Dirichlet and Neumann cases ($\gamma = \infty$) or ($\gamma = 0$) are known only in the limit cases $k = 0$ or $k = \infty$ (see [5]).

In particulary these inequalities solve the question if for certain $a+ib, b > 0$ and k there exists a sequence of smooth obstacles with uniformly bounded area such that $\sigma_{a,b}$ tends to infinity.

1.1 Proofs

Let u be the field of the outgoing wave satisfying (1),(2),(3). Everywere $\|\cdot\| = \|\cdot\|_{L_2(\partial\Omega,dS)}$. Lets prove that

$$\left\|\frac{\partial u}{\partial n} + (a+ib)u\right\| \ge b\|u\|. \tag{7}$$

Note that

$$\left\|\frac{\partial u}{\partial n}+(a+ib)u\right\|^2=\left\|\frac{\partial u}{\partial n}+au\right\|^2+2bIm\left(\int_{\partial\Omega}\frac{\partial u}{\partial n}\overline{u}dS\right)+b^2\|u\|^2.$$

Using the well known fact (evidence from the Second Green's identity)

$$Im\left(\int_{\partial\Omega}\frac{\partial u}{\partial n}\overline{u}dS\right)=k\sigma_{a,b}\geq 0, \tag{8}$$

we obtain (7). Note now that from (3) follows that

$$\left\|\frac{\partial u}{\partial n}+(a+ib)u\right\|=\left\|\frac{\partial e^{ikz}}{\partial n}+(a+ib)e^{ikz}\right\|\leq$$

$$\left\|\frac{\partial e^{ikz}}{\partial n}\right\|+\sqrt{a^2+b^2}\|e^{ikz}\|\leq\sqrt{S}(k+\sqrt{a^2+b^2})$$

So using (7) we obtain

$$b\|u\|\leq\sqrt{S}(k+\sqrt{a^2+b^2}), \tag{9}$$

and, in particularly, (6). Also from (3) we have for $r\in\partial\Omega$

$$-\frac{\partial u}{\partial n}=(a+ib)u+\frac{\partial e^{ikz}}{\partial n}+(a+ib)e^{ikz},$$

therefore

$$\left\|\frac{\partial u}{\partial n}\right\|\leq\sqrt{a^2+b^2}\|u|+\left\|\frac{\partial e^{ikz}}{\partial n}\right\|+\sqrt{a^2+b^2}\|e^{ikz}\|\leq$$

$$\sqrt{a^2+b^2}\|u\|+\sqrt{S}(k+\sqrt{a^2+b^2})\leq\frac{\sqrt{S}(k+\sqrt{a^2+b^2})(b+\sqrt{a^2+b^2})}{b}.$$

Now from (8) we have

$$\sigma_{a,b}\leq\frac{1}{k}\|u\|\|\frac{\partial u}{\partial n}\|\leq\frac{1}{k}\left(\frac{\sqrt{S}(k+\sqrt{a^2+b^2})}{b}\right).$$

$$\left(\frac{\sqrt{S}(k+\sqrt{a^2+b^2})(b+\sqrt{a^2+b^2})}{b}\right)=\frac{S(k+\sqrt{a^2+b^2})^2(b+\sqrt{a^2+b^2})}{kb^2}$$

Theorem 1 is proved.

References

[1] A.I.Aleksenko, W. de Roeck, E.L.Lakshtanov,“Resistance of the Sphere to a Flow of Quantum Particles”, J.Phys. A.Math.Gen, (39), pp. 4251-4255, 2005.

[2] G.A.Kriegsmann, “Alternate proof of the optical theorem for impenetrable targets”, *J Acoust Soc Am.*, Jan;119(1):31-2 (2006)

[3] A. Majda, M.E.Taylor, “The asymptotic behavior of the diffractive peak in classical scattering”, *Comm. pure and applied math.* vol. XXX, 639–669, (1977)

[4] A.G.Ramm, “Scattering by Obstacles” (Dordrecht: Reidel), (1986)

[5] W. De Roeck, E.L.Lakshtanov, “Total cross section exceeds transport cross section for quantum scattering from hard bodies at low and high wave numbers”, *J.Math.Phys*, 48, (2007).

[6] A.N.Tychonov, A.A.Samarsky, “Equations of Mathematical Physics Pergamon”, Oxford, (1963).